Background of the study

Pre-licensure baccalaureate nursing students undergo a significant amount of stress throughout a nursing program. However, pre-licensure baccalaureate nursing students receive education on the theoretical content on self-care practices for patients but oftentimes do not utilize these practices to manage their own anxieties. Self-care practices are typically taught throughout the baccalaureate curricula. According to the Essentials of Baccalaureate Education for Professional Nursing Practice, the baccalaureate nursing graduate should be able to provide self-care instruction to others (American Association of Colleges of Nursing [AACN], 2008). Nursing students learn the application of self-care practice education for their patients and communities. However, many nursing students lack experience with personal self-care and health promotion (Nevins & Sherman, 2016), and may not be applying the practice of positive self-care strategies to help manage their own anxiety (Ashcraft & Gatto, 2015). In a recent study, nursing students who participated in a self-care course identified the need to provide self-care for themselves, learned about the application in their personal and professional lives, felt empowered to educate and advocate for their patients, and recognized the need to retain the knowledge for the future (Padykula, 2017). Additional studies have stated that nursing students who engage in self-care activities enjoy learning these practices (Blum, 2014), report both lower levels of stress (Drew et al., 2016), and lower levels of anxiety (Kang, Choi, & Ryu, 2009). Even with this knowledge, nursing students are at-risk for developing maladaptive coping strategies to stress. In fact, Drew et al. (2016) concluded that nursing students engage in less self-care practices as they progress through a nursing program and recommended that self-care practices be reinforced throughout the nursing curriculum.

Nursing students have reported that insufficient information, money, and rest have limited their ability to engage in positive health-promoting behaviors (Bryer, Cherkis, & Raman, 2013). Baccalaureate nursing students report that stress, anxiety, and time commitments are barriers to self-care behaviors in nursing school, even though they would like to improve their well-being with adequate sleep, exercise, diet, and hydration (Nevins & Sherman, 2016).

Unfortunately, nursing students who are unable to manage their anxiety may develop low levels of self-confidence which can negatively impact their ability to learn. Anxiety can impair learning which leads to poor performance in the clinical environment (Cheung & Au, 2011; Melincavage, 2011; Melo et al., 2010; Nielsen & Harder, 2013), decrease levels of self-confidence (DeBrew & Lewallen, 2014), and contribute to low levels of self-efficacy (Watt et al., 2016). Additionally, nursing students who are unable to manage their anxiety are at-risk for clinical failure, hinder faculty-student and patient-student interactions, and present themselves as unprofessional and unsafe (DeBrew & Lewallen, 2014; Killiam, Luhanga, & Bakker, 2011). These negative outcomes can impact students can affect students throughout a nursing program

reduce anxiety and improve the health-promoting behaviors of interpersonal support and stress management.

References

American Association of Colleges of Nursing. (2008). The essentials of baccalaureate

education for professional nursing practice. Retrieved form http://www.aacn.nche'

Rababah, J. (2016). Care for the caregiver: Evaluation of mind-body self-care for

accelerated nursing students. Holistic Nursing Practice, 30(3), 148-154.

doi:10.1097/HNP.0000000000000140

Kang, Y. S., Choi, S. Y., & Ryu, E. (2009). The effectiveness of a stress coping program

based on mindfulness meditation on the stress, anxiety, and depression

experienced by nursing students in Korea. Nurse Education Today, 29(5), 538-

543. doi:10.1016/j.nedt.2008.12.003

Killiam, L. A., Luhanga, F., & Bakker, D. (2011). Characteristics of unsafe undergraduate nursing students in clinical practice: An integrative literature review. Journal of Nursing Education, 50(8), 437-446. doi:10.3928/01484834- 20110517-05

Journal of Nursing Management, 19(5), 572-584. doi:10.1111/j.1365-

2834.2011.01239.x

Watt, E., Murphy, M., MacDonald, L., Pascoe, E., Storen, H., & Scanlon, A. (2016). An

evaluation of a structured learning program as a component of the c0 a65

Protected population and sensitive subjects: Indicate if any Human Subjects from the following list will be involved in the proposed activity:

1R D QZ H USURY GH G

Vulnerable Subjects

No vulnerable subjects will be used in this study.

If it is your intention to use vulnerable subjects, justify the importance of their use. Here and throughout

Method of Subject Selection

The setting for this study will be up to six public or private universities located in the Mid-Atlantic region. All potential schools of nursing have a traditional four-year baccalaureate degree in nursing program. The potential schools were identified through a review of the NCLEX-RN Performance of First Time Candidates Educated in Pennsylvania who Completed NCLEX in the U.S. published by the National Council of State Boards of Nursing. Nursing programs who had at least 50 graduates who were eligible to take the NCLEX in 2017 will be recruited. In Pennsylvania, 24 schools of nursing are eligible to be recruited based on these requirements. From the 24 schools of nursing in Pennsylvania, eight nursing schools were excluded because they have a PhD program. Pennsylvania State University has multiple campuses that offer traditional BSN programs. Two of the campuses will be excluded because they offer PhD programs (University Park & Hershey). However, this leaves five satelitte campuses that may be recruited from. If none of these 16 schools of nursing are agreeable to participate, the researcher will look to other traditional baccalaureate nursing schools in the Mid-Atlantic region.

An email will be sent to the nursing chairs or faculty contact at each of the 16 eligible schools of nursing in Pennsylvania. The email will describe the research study purpose, timeline, and ask if the nursing chair or faculty contact would be interested in being a study site. If the nursing chair or faculty contact is agreeable, an in-person or telephone meeting will be offered to discuss the project. The first six schools of nursing who are agreeable to participate will be used for the study. Site approval or site IRB approval will be completed at each instituion.

The researcher will go to each school in-person, provide a Zoom session, or offer to pre-record a video to introduce the study and provide the initial link to the pre-licensure senior-level baccalaureate nursing students. If the school would rather have the nursing secretary or site designee at the school distribute an initial email, the researcher will provide a precorded video to explain the study. Student recruitment will start at the beginning of the Spring semester of 2018.

A power analysis was conducted to estimate the sample size required for the quantitative data analysis. According to Polit and Beck (2012), many nursing research studies utilize an effect size between .20 to .40 to determine the relationship between independent and

3 UR Y GH RP SOH W QL RU D WR D EXWKRZ UH YH DFN VXEM HW F ZLOO EH GH QL H G H EXLW G LQY W GW R SD UF LSD W IF WQGLFD W D SSUR[LP D W O KRZ P D Q UH YH DFN VXEM HW \ RX ZLOO FRQ W D F Q KRZ P D Q \ RX ZLOO D FX D O XVH Q \ RXU UH YH DFN < RXU GH Y FULSWR RI H EXLW H Q D Q \H H EXLW P H Q D Q VH H F RQ P XW Q F XGH D\Q GH W W U D Q RXQF H PH Q V DY GH W H PH Q V RU RWKH UUH D W GP D W O D W O D \ $Q\ P D W O D \ W

dependent variables within the study population. Therefore, a medium effect size of .30 will be used. The standard significance level of 5% and .80 for power will be used (Polit & Beck, 2012). Based on an effect size of .30, significance level of 5%, and a power of .80, a sample size of 72 study participants was computed using G*Power Version 3.1.9.2. Therefore, it is anticipated that nursing students from up to six public or private institutions in the Mid-Atlantic region will be recruited. The attrition of study subjects needs to be addressed due to the three data collection time points. The attrition of study subjects is common in a longitudinal study and is rarely 0% (Polit & Beck, 2012). Therefore, an attrition rate of 20% was computed for this study indicating that 14 additional study participants, for a total of 86 study participants, would need to be recruited to overcome anticipated attrition or ruled

Study Site

I will be seeking approval from the IRB at BSN programs in the Mid-Atlantic Region. These are the potential study sites. I plan to have up to 6 study sites. I plan to use personal and professional contacts to connect with faculty members or nursing chairs at these institutions.

Potential study sites
Bloomsburg University BSN
Cedar Crest College BSN
DeSales University BSN
Edinboro University BSN
Holy Family University BSN
Jeffers

Potential Benefits

There may be no direct benefits from participation in the study. However, the knowledge received may be v

researcher will receive a Microsoft® Excel document that will contain no personally identifiable information of nursing students from the ARL. The participants will provide a ten-digit cell phone number and email address over the c

Will the research be conducted in established or commonly accepted educational settings, involving normal educational practices, such as (i) research on regular and special education instructional strategies, or (ii) research on the effectiveness of or the comparison among instructional techniques, curricula, or classroom management methods?

Yes

Will the research be involving the use of educational tests (cognitive, diagnostic, aptitude, achievement), survey procedures, interview procedures or observation of public behavior, unless: (i) information obtained is recorded in such a manner that human subjects can be identified, directly or through identifiers linked to the subjects; and (ii) any disclosure of the human subjects' responses outside the research could reasonably place the subjects at risk of criminal or civil liability or be damaging to the subjects' financial standing, employability, or reputation.

No

Will the research be involving the use of educational tests (cognitive, diagnostic,

Expedited Instructions

,Q \RXU MGXJPHQW GRHV\RXU SURMHFW DOOXQGHU RQH RI WKH FDWHJRULHV VHOHFWHG IRU H[SHGLWHG UHYLHZ DVVXPH GEHRZ " ,I \RX EHOLHY HWGRHV QGLFD WKH HDWJRU R ZKLFK \RXU FODLPLQJ H[SHGLWHG UHYLHZ EH FKRRVLQJ \HVQH[WWR WKH HUHOY DVQDD WJRU

Clinical studies of drugs and medical devices only when condition (a) or (b) is met. a. Research on drugs for which an investigational new drug application (21 CFR Part 312) is not required. (Note: Research on marketed drugs that significantly increases the risks or decreases the acceptability of the risks associated with the use of the product is not eligible for expedited review.) b. Research on medical devices for which (i) an investigational device exemption application (21 CFR Part 812) is not required; or (ii) the medical device is cleared/approved for marketing and the medical device is being used in accordance with its cleared/approved labeling.

No

Collection of blood samples by finger stick, heel stick, ear stick, or venipuncture as follows: a. from healthy, nonpregnant adults who weigh at least 110 pounds. For these subjects, the amounts drawn may not exceed 550 ml in an 8 week period and collection may not occur more frequently than 2 times per week; or b.from other adults and children2, considering the age, weight, and health of the subjects, the collection procedure, the amount of blood to be collected, and the frequency with which it will be collected. For these subjects, the amount drawn may not exceed the lesser